




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



The first pregnant woman with COVID-19 in Venezuela: Pre-symptomatic transmission

Dear editor

Since the first confirmed case from Latin America in February 25th, 2020 in São Paulo, Brazil, SARS-CoV-2 cases have rapidly began to spread across the Latin America [1]. Available data about COVID-19 and its impact in this region is limited, including pregnant women data. Here we present the first confirmed case of a pregnant woman with COVID-19 in Venezuela, who was presumed to be infected by her mother during her pre-symptomatic phase.

On March 28th, a 32-year-old female, with 38-week of pregnancy, without comorbidities, presented to the University Hospital of Caracas, with dry cough, and fatigue. Her symptoms started 12 days prior, when she noticed headache and anosmia. She had no previous travel history or contact with any respiratory-symptomatic person before onset of symptoms. On admission, physical examination and laboratory findings revealed no relevant alterations.

The patient consulted accompanied by her mother, who had been in the United States since October/2019, followed by a trip to Peru on February 22nd where she remained in Lima until March 10th, when she returned to Caracas. The mother was asymptomatic until March 25th

(Fig. 1). The pregnant patient and her mother lived together. Nasopharyngeal swab samples were collected from both which tested positive for SARS-CoV-2 real-time RT-PCR at the “Rafael Rangel National Hygiene Institute”. The patient’s husband and her two other children were asymptomatic, nasopharyngeal swab samples were collected for the complete family nucleus; testing negative for SARS-CoV-2. On March 31st, 2020, she was admitted for a caesarean section. Her son was born without complications, testing negative for SARS-CoV-2 (rRT-PCR).

SARS-CoV infection has been associated with poor pregnancy outcomes, including critical maternal illness [2]. Similarly, SARS-CoV-2 associated preterm birth has been reported [3]. This case’s presentation showed no significant alterations related to COVID-19, as it has been recently reported in a retrospective study [4]. On the other hand, to date, there has been no evidence of vertical transmission of COVID-19. In this case, three nasopharyngeal swabs obtained from the newborn infant were repeatedly negative for SARS-CoV-2.

Interestingly, the pregnant woman developed symptoms before her contact did. This could be explained in part, by the immunosuppressed state of pregnancy where there is attenuation of the cell-mediated immunity by Th1 cells due to the physiological shift to a Th2 dominant




Fig. 1. Timeline of the pregnant patient: symptoms and results of rtPCR testing for SARS-CoV-2 by date.

<https://doi.org/10.1016/j.tmaid.2020.101805>

Received 21 April 2020; Received in revised form 7 May 2020; Accepted 19 June 2020

Available online 25 June 2020

1477-8939/© 2020 Elsevier Ltd. All rights reserved.

environment [5], with this contributing to overall infectious morbidity by increasing maternal susceptibility to intracellular pathogens like viruses. Additionally, person-to-person transmission within cluster families, by pre-symptomatic [6], or even asymptomatic infectors has been previously reported.

With the increase of the pandemic in Latin America, a region with high fecundity rates and a fragile health system we should expect to see more cases of COVID-19 among pregnant women. Therefore, pregnant women and their newborn babies should be considered a key high-risk population in the development of prevention and management strategies for COVID-19. This vulnerable group needs to be studied in detail to better understand its clinical impact.

Declaration of competing interest

All authors declare that they have no conflicts of interest.

References

- [1] Rodriguez-Morales Alfonso J, et al. COVID-19 in Latin America: the implications of the first confirmed case in Brazil. *Trav Med Infect Dis* 2020;101613. <https://doi.org/10.1016/j.tmaid.2020.101613>.
- [2] Wong Shell F, et al. Pregnancy and perinatal outcomes of women with severe acute respiratory syndrome. *Am J Obstet Gynecol* 2004;191:292–7. <https://doi.org/10.1016/j.ajog.2003.11.019>.
- [3] Wangla Xiaotong, et al. A case of 2019 Novel Coronavirus in a pregnant woman with preterm delivery. *Clin Infect Dis* 2020. <https://doi.org/10.1093/cid/ciaa200>.
- [4] Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet* 2020;395:809–15. [https://doi.org/10.1016/S0140-6736\(20\)30360-3](https://doi.org/10.1016/S0140-6736(20)30360-3).
- [5] Saito S, et al. Distribution of Th1, Th2, and Th0 and the Th1/Th2 cell ratios in human peripheral and endometrial T cells. *Am J Reprod Immunol* 1999;42:240–5. <https://doi.org/10.1111/j.1600-0897.1999.tb00097.x>.
- [6] Qian Guoqing, et al. A COVID-19 Transmission within a family cluster by pre-symptomatic infectors in China. *Clin Infect Dis* 2020. <https://doi.org/10.1093/cid/ciaa316>.

David A. Forero-Peña *

*Infectious Diseases Department of the University Hospital of Caracas,
Caracas, Venezuela
Biomedical Research and Therapeutic Vaccines Institute, Ciudad Bolívar,
Venezuela*

Mariana I. Rodríguez, David M. Flora-Noda, Andrea L. Maricuto, Viledy L. Velásquez, Lili M. Soto, Yoel D. García, Francisco A. Ureña, Mariangela Mosqueda, Rommel Franco-Lugo
*Infectious Diseases Department of the University Hospital of Caracas,
Caracas, Venezuela*

María V. Marcano
*Biomedical Research and Therapeutic Vaccines Institute, Ciudad Bolívar,
Venezuela*

Grismely A. Morillo, Martin Carballo, Jocays Caldera, María Carolyn Redondo, María Eugenia Landaeta
*Infectious Diseases Department of the University Hospital of Caracas,
Caracas, Venezuela*

* Corresponding author. Infectious Diseases Department of the University Hospital of Caracas, Caracas, Chaguaramos parish of Libertador Municipality in the District Capital, Venezuela. Tel.: +5804249548706.

E-mail address: vacter.cv@gmail.com (D.A. Forero-Peña).